

Subject index

| | | | |
|-------------------------------|----------|---------------------------|----------------|
| A | | E | |
| Academic yield | 198 | <i>Eisenia andrei</i> | 320 |
| Acclimatization | 216 | <i>Elaeis guineensis</i> | 320 |
| All possible regressions | 198 | <i>Elaeis guineensis</i> | 421 |
| <i>Aloe vera</i> L. | 216 | Electrical conductivity | 396 |
| Amaranth | 115 | Endophytic fungi | 274 |
| <i>Amaranthus cruentus</i> | 115 | Establishment | 216 |
| Ammonium | 71 | Experimental error | |
| <i>Anacardium occidentale</i> | 18 | variance | 97 |
| <i>Annona muricata</i> L. | 2 | Extraction | 302, 385 |
| <i>Atta</i> | 311 | extraction methods | 472 |
| Axenic | 29 | | |
| <i>Azolla</i> , | 136 | F | |
| | | Fabaceas | 385 |
| B | | Fertility | 332 |
| Banana | 430 | FHIA varieties | 430 |
| Biofertilizer | 136 | Flowering | 162 |
| Biologic control | 45 | Freezing | 235 |
| Black Sigatoka | 295 | Fructification | 162 |
| Budding | 2 | | |
| Buffelgrass | 151 | G | |
| | | Géneros | 225 |
| C | | germination | 18, 189, 258 |
| Carotenoids | 302 | Gramineous | 332 |
| <i>Cenchrus ciliaris</i> | 151 | Grape | 460 |
| Cereals | 115 | Grassland management | 339 |
| Charcoal root | 286 | Growing rate | 136 |
| <i>Chlorella sorokiniana</i> | 29 | growth | 2, 61, 71, 408 |
| Clotted | 162 | Guava | 61 |
| Cocoa | 350 | Guinea grass | 85 |
| Coconut coir dust | 189 | | |
| Color | 484 | H | |
| Comparison of designs | 97 | Humidity | 286 |
| Compost | 189 | | |
| Corn | 374 | I | |
| corn bread | 235 | Incidence | 61, 182 |
| Cp of Mallows | 198 | Instant beverage | 115 |
| Cultivars | 408, 449 | Interference | 374 |
| Dietetic fiber | 235 | Intraspecific competition | 151 |
| Distribution | 286 | Irrigation | 61 |
| <i>Dothiorela</i> sp. | 61 | | |
| Doubling time | 136 | K | |
| Drying storage | 350 | Koch's postulates | 421 |
| Dummy Variables, | 198 | | |
| Dynamic | 85 | | |

| | | | |
|--------------------------------|----------|--|----------|
| L | | Pollen germinability | 173 |
| Leaching | 396 | pollen quantity | 173 |
| Leaf-cutting ants | 311 | pollen viability | 173 |
| Legumes | 332 | Post-harvest fermentation | 350 |
| <i>Lemna</i> sp. | 71 | Potassium | 162, 484 |
| <i>Leucaena leucocephala</i> | 258 | Press | 198 |
| | | <i>Prosopis juliflora</i> | 258 |
| M | | Q | |
| Macroelements | 162 | Qualitative and quantitative variables | 198 |
| <i>Macrophomina phaseolina</i> | 45 | Quality | 52, 430 |
| <i>Mangiferae indica</i> | 274 | | |
| <i>Manihot esculenta</i> | 311 | R | |
| Maracaibo | 151 | R ² | 198 |
| Melon | 449 | Regression coefficients variance | 97 |
| Model comparisons | 198 | Relative efficiency | 97 |
| Morphological characteristics | 18 | Repeated Incomplete Blocks, | 97 |
| Morphological characteristics | 258 | Resistance | 295 |
| MSPD | 472 | Rice bran | 235 |
| Multiplication | 216 | Ripening | 430 |
| | | Roots | 460 |
| N | | Rotary Central | |
| Nitrate | 71 | Compound design | 97 |
| Nitrogen | 85 | <i>Rottboellia cochinchinensis</i> | 374 |
| nitrogen fixation | 136 | | |
| Non-axenic | 29 | S | |
| | | Salinity | 29 |
| O | | Salinity | 396 |
| Oranges | 302 | Seasonal growth | 460 |
| Organic solvents | 302 | Seedling | 18 |
| Orthogonal Central | | Seedlings | 258 |
| Compound design | 97 | Sensorial analysis | 52 |
| | | <i>Sesamum indicum</i> | 45 |
| P | | Severity | 182 |
| Papaya | 173 | Soil series | 396 |
| Passalora <i>koepkei</i> | 182 | Soils | 332 |
| Pasture productions | 339 | Stepwise | 198 |
| Pestalotiopsis | 421 | Stomatal density | 295 |
| Pesticides | 472 | Strategies for pastures utilization | 339 |
| pH | 29 | Substrate | 189 |
| pH | 71 | Sugar cane | 182 |
| Phenology | 2 | | |
| Phosphorous | 136 | T | |
| Physical quality | 484 | Temperature | 29, 286 |
| Phytopathogens | 274 | <i>Theobroma cacao</i> L. | 52 |
| Plant spacing | 449 | | |
| Plantain, | 295 | | |
| Poaceae | 151, 225 | | |

| | | | |
|---------------------------|---------------|------------------|--------------|
| <i>Theobroma cacao</i> L. | 350 | W | |
| Tomato | 162, 189, 484 | Wastes | 320 |
| Transmission seed born | 385 | Weed | 151, 374 |
| U | | Y | |
| Urea | 71 | Yellow leaf spot | 182 |
| Vegetables | 472 | Yield | 61, 311, 449 |
| Venezuela | 151, 225 | Z | |
| Virus | 385 | Zulia | 225 |
| <i>Vitis sp.</i> | 460 | | |
| <i>Vitis vinifera</i> | 408 | | |